



GenX Exposure Study: PFAS blood levels in the Lower Cape Fear Region, NC 2020 to 2023

Jane Hoppin, ScD Professor, NC State Principal Investigator

October 29th, 2024







Agenda

Welcome
Introduction to PFAS and GenX Exposure Study
Overall study findings for PFAS blood levels
Overview of results for Lower Cape Fear Region
Future Plans
Q&A Panel



GenX Exposure Study Mission

The purpose of the GenX Exposure Study is to understand how PFAS chemicals in drinking water may impact human health of residents of the Cape Fear River Basin.



Tonight's Topic

PFAS levels in blood samples collected in 2023.





Background

PFAS, and PFAS in the Cape Fear River Basin, & The GenX Study



Per- and Polyfluoroalkyl Substances (PFAS)

Resistant to:

Water

Stains

UV radiation

Used in consumer products since the 1950s:

Surfactants, lubricants, adhesives

Carpet, upholstery, clothing

Car interiors

Food packaging, nonstick cookware

Cleaning products

Personal care products

Fire-fighting foam

Chemical Properties Lead to:

Persistence and bioaccumulation

Ubiquitous in indoor environment and blood (e.g., NHANES)

Transport around globe in ocean currents and atmosphere





Cape Fear River Basin, North Carolina Haw River Largest watershed in NC Deep River Supplies ~1.5M people Cape Fear River with drinking water fluorochemical manufacturing facility

GenX Study Background: Exposure Study 2017-2019

In 2017, we started in Wilmington, NC.

In 2019, we included the Fayetteville private well community.

- Found high levels of PFAS in people's blood.
- Identified new PFAS associated with the Chemours.
- Nafion byproduct 2, PFO5DoA, PFO4DoA in almost everyone from
- Wilmington.
- Nafion byproduct 2 and PFO5DoA in some people from Fayetteville.
- We did not find GenX in people's blood.



GenX Cohort Health Study: 2020 – date

In 2020-21, we enrolled 1020 people throughout the Cape Fear River Basin to measure PFAS exposure in blood and to follow them for up to 20 years to learn about health effects.

We included Pittsboro to characterize PFAS upriver from the chemical plant.

In 2023, we resampled 519 people and enrolled 72 new people to improve the representation of people in the sample.

In 2024, we are resampling the people who did not participate in 2023.



Study Design: GenX Cohort Study

Enroll people ages 6 and older from 3 regions in the Cape Fear Basin

Lower Cape Fear

Fayetteville

Pittsboro

Started in 2020-21

Enrolled >1000 people

Resampled in 2023

Plan to follow people for up to 20 years

Collect blood

Analyze for GenX and other PFAS

Analyze for lipids, thyroid hormones, and comprehensive metabolic panel

Report back results to community, individuals






Who is in the study?

People range in age from 6 to 92 years old.

Most people are 60 and older

About 55% female





Sample Collection: Summer-Fall 2023

Blood sample

Clinical measures

- lipids,
- comprehensive metabolic panel,
- thyroid hormones
- Results returned in December 2023

Serum PFAS

- 41 analytes in 2023
- Results returned in April-May 2024

Body composition

Questionnaire

Update health history





Sample Collection 2023

Collected blood from 591 people

118 Pittsboro

126 Fayetteville

347 Lower Cape Fear

Resampled 523 of original 1019 (~50%)

Enrolled 72 new people

Mostly in Lower Cape Fear Region



Two Letters to Study Participants about 2023 results

Individual PFAS Results

Your individual PFAS values for 2023 and 2020/21 Your individual NASEM recommendations April-May 2024

Community PFAS results

Shows where your PFAS results fit in your community July 2024



Key Overall Findings

We measured blood samples from over 500 people for 41 different PFAS. On average, levels of PFAS in blood are lower in 2023 than 2020-21

Not all people's blood levels went down

PFAS blood levels are still higher than most people in the US

People throughout the Cape Fear River Basin from Pittsboro to New Hanover and Brunswick Counties are adversely impacted by a variety of PFAS.



Sampling in 2020-2021

Collected blood from 1,019 people across the entire study area

Lower Cape Fear Region

Nov 2020: 282 people

Nov 2021: 300 people

We tested for 44 different PFAS

Sampling in 2023

Collected blood from 591 people across the entire study area

Lower Cape Fear Region

347 total

64 new people

We tested for 41 different PFAS



Tested blood samples for 41 PFAS in 2023

7 PFAS commonly found in people			
	PFOS		
	PFOA		
	PFHxS		
	PFNA		
	PFDA		
	PFUnDA		
	MeFOSAA		
Other perfluoroalkyl carboxylic acids			
•	PFBA		
	PFPeA		
	PFHxA		
	PFHpA		
	PFTrDA		
	PFDoA		
	PFTeDA		
	PFHxDA		
	PFODA		
Other perfluoroa	lkyl sulfonic acids		
	PFBS		
	PFPeS		
	PFHpS		
	PFNS		
	PFDS		

Perfluoroether carboxylic acids		
	PEPA	
	GenX	
	PFO3OA	
	PFO4DA	
	PFO5DoA	
	NaDONA	
Perfluoroether sulfonic acids		
	Nafion byproduct 1	
	Nafion byproduct 2	
	F53B Major (9CI-PF3ONS)	
Perfluoroalkyl sulfonamides		
	6:2 FTS	
	FHxSA	
	FOSA	
	MeFOSA	
	F53B Minor	
	(11CI-PF3OUdS)	
	FBSA	
	NEtFOSAA	
Fluorotelomer carboxylic acid		
	7:3 FTCA	

Fluorotelomer sulfonic acids		
	10:2 FTS	
	8:2 FTS	
	4:2 FTS	

Samples analyzed at Eurofins laboratory in Sacramento, CA



Focus today on these PFAS

7 PFAS commonly	found in people
	PFOS
	PFOA
	PFHxS
	PFNA
	PFDA
	PFUnDA
	MeFOSAA
Other perfluoroalkyl carboxylic acids	
	PFBA
	PFPeA
	PFHxA
	PFHpA
	PFTrDA
	PFDoA
	PFTeDA
	PFHxDA
	PFODA
Other perfluoroall	kyl sulfonic acids
	PFBS
	PFPeS
	PFHpS
	PFNS
	PFDS

Perfluoroether carboxylic acids		
	PEPA	
	GenX	
	PFO3OA	
	PFO4DA	
	PFO5DoA	
	NaDONA	
Perfluoroether sulfonic acids		
	Nafion byproduct 1	
	Nafion byproduct 2	
	F53B Major (9CI-PF3ONS)	
Perfluoroalkyl sulfonamides		
	6:2 FTS	
	FHxSA	
	FOSA	
	MeFOSA	
	F53B Minor	
	(11CI-PF3OUdS)	
	FBSA	
	NEtFOSAA	
Fluorotelomer carboxylic acid		
	7:3 FTCA	

Fluorotelomer sulfonic		
acids		
	10:2 FTS	
	8:2 FTS	
	4:2 FTS	



Chemours-related PFAS

What about the Chemours-related PFAS

In 2023, we continued to find PFO5DoA and Nafion byproduct 2 in the blood of most people from the Lower Cape Fear Region & about 20% of the people from Fayetteville.

Overall, the levels continue to decrease over time.

	Lower Cape Fear Region	Fayetteville
PFO5DoA	90%	18%
Nafion byproduct 2	57%	21%



Change in median levels in blood samples (ng/mL) since 2020

Nafion byproduct 2 Change in Median PFO5DoA concentrations in LCFR 2020 2021 2023 Change in Median Nafion byproduct 2 concentrations in LCFR 2020 2021 2023 2020 0.5 2020 2021 Concentration (ng/mL) Concentration (ng/mL) 2021 2023 2023 0.1 0.0 Nafion byproduct 2 PFO5DoA



Four PFAS were found in almost everyone:

PFOS

PFOA

PFHxS

PFNA




Strip Charts of Individual PFAS Across the Lower Cape Fear Region Samples 2023






N = 347

Stripchart for PFOS for entire study, 2023






Stripchart for PFOS for the Lower Cape Fear Region, 2023






Lower Cape Fear Region 2023 PFOS Stripchart to Boxplot






Lower Cape Fear Region 2023 PFOS Stripchart to Boxplot






PFAS Levels for 4 most common PFAS in the Lower Cape Fear Region 2023





N = 347


PFOS & PFOA Levels in 2023 by community compared to US population for 2017-2020





2023 GenX Samples

PFHxS and PFNA Levels from 2023 by community compared to US population for 2017-2020





2023 GenX Samples

How have PFAS blood levels changed over time?

For the common PFAS (PFOS, PFOA, PFHxS, PFNA), the levels in blood have decreased over time

In people with two measurements (N=519)

PFOS dropped 1.7 ng/mL (24%)

PFOA dropped 1 ng/mL (32%)


PFHxS dropped 0.5 ng/mL (21%)

In a 2-3 year period

Results are similar over all communities.



PFAS Blood Levels in 2020 vs 2023 in the Lower Cape Fear Region samples







Not all people's PFAS levels went down

We measured 41 PFAS in people's blood

Some individuals had one or more PFAS increase

We are working to understand why?

Maybe a different exposure source than water

Food, cleaning products, occupational exposures?

Change in body composition

We're measuring chemicals in people's bodies, so changes in your body may affect the concentration we measure

Analytical variability

Some levels are really low and may bounce around a bit.

Reach out to us if you want help identifying why your levels changed.



Key Findings


We measured blood samples from over 500 people for 41 different PFAS.

- 1. On average, levels of PFAS in blood are lower in 2023 than 2020-21
- 2. Not all people's blood levels went down
- 3. PFAS blood levels are still higher than most people in the US
- 4. People throughout the Cape Fear River Basin from Pittsboro to New Hanover and Brunswick Counties are adversely impacted by a variety of PFAS.




What can I do with this information?

NASEM Recommendations, Study Retention, and Exposure Control



NASEM 7 in 2023



Sum of 7 serum PFAS (PFOS, PFOA, PFHxS, PFNA, PFDA, MeFOSAA, PFuNDA)

Does not include PFHps, Nafion byproduct 2, PFO5DoA

Overall ~20% exceed 20 ng/mL

The fact that levels are now lower doesn't negate the fact that levels were higher.

≥ 20 ng/mL summed PFAS

Higher risk of adverse effects
Reduce exposure
Also test for thyroid function, kidney and testicular cancer, ulcerative colitis

2 - <20 ng/mL summed PFAS

Potential for adverse effects in sensitive populations Reduce PFAS exposure Screen for dyslipidemia, hypertensive disorders of pregnancy, and breast cancer

< 2 ng/mL summed PFAS

Adverse health effects not expected. Recommend usual standard of care.



What can you do about PFAS exposure and health effects?

If you're in the study, please stay in the study. This lets us learn how the chemicals are moving through our bodies and potential health effects.

If your PFAS levels have increased, think about potential ways you might come into contact with PFAS

Home grown fruits and vegetables, eggs

Occupational Exposures

and consider potential changes in your body.

Weight change

New medication



What can you do about PFAS exposure and health effects?

If you live in a region in the Cape Fear River, consider your potential for PFAS exposure. Because almost all people in the study were classified at some potential adverse risk, talk to your health care provider about actions you can take to protect your health.

Think about how you may be exposed to PFAS.

Home grown fruits and vegetables, eggs

Occupational Exposures



If you want to test for PFAS in your blood:

There are resources on the GenX Study website to help you find the testing you want.



What is next?

Resampling and Scientific Papers



What's next for study? 2024

We are resampling anyone who enrolled between 2020 and 2023 to ensure that we have two measurements per person.

If we missed you last year, we will be contacting you to make sure we include you this year.

If you've moved, changed your email or phone, please let us know. genx-exposure-study@ncsu.edu or call us at (855) 854-2641

We'll be sampling those we missed in 2023 in Wilmington on Nov 8 and 9, 2024.

You can schedule with us at the end of the meeting.



What's next for the Study: Health effects evaluation

We are currently working on scientific papers looking at how PFAS may affect
Thyroid hormones
Liver enzymes

We hope to publish these by early 2025 and share these results with you. If you have ideas for things we should look at, please reach out.



If you want to learn more:

Check out our website (genxstudy.ncsu.edu)
Follow our Instagram! @ncsu_genx_study



Thank you to our funders and partners!

- NC State
- ECU
- Cape Fear River Watch
- Haw River Assembly
- Sustainable Sandhills
- New Hanover County Health Department
- Cumberland County Health Department
- Chatham County Health Department
- New Hanover County NAACP
- Warner Temple AME Zion Church
- Town of Navassa
- Our wonderful study participants

Funding: NIEHS R21 R21ES029353, P42ES031009

CHHE P30ES025128

Matching Funds from NC Policy Collaboratory



Welcome to our panelists!

Ralph Mead, PhD

PFAS Testing Network, Project Director

Rachylle Hart

PFAS Testing Network, Program Coordinator

Sanjay Batish, MD

Practicing at Batish Family Medicine in Leland, NC since 1998

Hanna Bailey, MPH

GenX Exposure Study, Moderator

